

FOCUSED ABDOMINAL ULTRASOUND: STILL RUNNING? PRINCIPLES AND PITFALLS

Ruth Løkke, MD

Definition

Focused Assessment with Sonography in Trauma (FAST) is a quickly performed screening technique aimed at exploring the deep peritoneal recesses to detect large collections of free fluid, as an indirect sign of acute hemorrhage and injury to visceral organs[1, 2].

Principles

FAST is performed as part of the 1. or 2. survey of the trauma patient in the admission area to determine whether immediate surgery is needed before the patient undergoes a further evaluation with CT.


The following four standard views are obtained with the patient in supine position:

Pericardial: Transverse view of the subxiphoid region to show pericardial effusion and the left lobe of the liver.

Perihepatic: Longitudinal view of the right upper quadrant to show the right lobe of the liver, the right kidney, and the space between the two (the Morison pouch).

Perisplenic: Longitudinal view of the left upper quadrant to show the left kidney, the spleen, and the space between them.

Pelvic: Transverse and longitudinal views of the suprapubic region to show the urinary bladder and retrovesical or rectouterine pouch, the pouch of Douglas.



A positive FAST is free fluid detected in one or more of the four views, whereas a negative FAST means no free fluid detected. For hemodynamically unstable patients, a positive FAST should be followed by laparotomy, whereas a negative result suggests a search for other sources of hemorrhage. FAST can easily be repeated in the unstable patient, when suspicion of intraperitoneal hemorrhage is high. For stable patients a positive FAST result suggests CT, whereas a negative result suggests observation, follow-up FAST or CT[3].

Pitfalls and limitations

FAST sensitivity has been reported ranging widely: 0.64–0.98[4-7], and is even more decreased in patients with pelvic fractures[8]. Specificity is high, at 0.86–1.00.

Pitfalls leading to false positive results[9]:

- Iatrogenic free fluid
- Pelvic physiologic fluid (female)
- Loops of fluid filled bowel

Pitfalls leading to false negative results[9]:

- Incomplete or empty bladder
- Echogenic clot
- Contained injury
- Small amounts of free fluid, volumes less than 400 ml[10, 11]

FAST results are also dependent on patient habitus and injury (e.g. subcutaneous edema) and operator experience. FAST is performed in the emergency unit with fully enlightened room, other simultaneous procedures, reduced space, portable equipment and short time of scanning, leading to decreased accuracy of the examination.

Despite scanning in the areas for the liver, spleen and kidneys, sonography in trauma settings has showed low sensitivity for organ-specific diagnoses[12]. Therefore, when performing FAST, the focus is exclusively on detection of free intraperitoneal fluid. Hypotensive patients with mesenteric or bowel injuries can present without free fluid.

Sonography in general is not able to show spinal and pelvic fractures or diaphragmatic ruptures and has limited accuracy for vascular, pancreatic, adrenal, bowel and mesenteric injuries.

Recommendations

FAST is recommended as a part of the initial evaluation of trauma patients. Only experienced radiologists or other physicians with high ultrasound work-load should perform the FAST. Standard scans should be quickly performed and stored, without delaying the transfer of the patient to CT or the operating room. A positive FAST in hemodynamically unstable patients should lead to immediate surgery.

Reference List

1. Korner M, Krotz MM, Degenhart C, Pfeifer KJ, Reiser MF, Linsenmaier U (2008) Current Role of Emergency US in Patients with Major Trauma. *Radiographics* 28:225-242.
2. Catalano O, Siani A (2004) Focused assessment with sonography for trauma (FAST): what it is, how it is carried out, and why we disagree. *Radiol Med (Torino)* 108:443-453.
3. Scalea TM, Rodriguez A, Chiu WC, Brenneman FD, Fallon WF, Jr., Kato K, McKenney MG, Nerlich ML, Ochsner MG, Yoshii H (1999) Focused Assessment with Sonography for Trauma (FAST): results from an international consensus conference. *J Trauma* 46:466-472.
4. McGahan JP, Rose J, Coates TL, Wisner DH, Newberry P (1997) Use of ultrasonography in the patient with acute abdominal trauma. *J Ultrasound Med* 16:653-662.
5. Forster R, Pillasch J, Zielke A, Malewski U, Rothmund M (1993) Ultrasonography in blunt abdominal trauma: influence of the investigators' experience. *J Trauma* 34:264-269.
6. Nunes LW, Simmons S, Hallowell MJ, Kinback R, Trooskin S, Kozar R (2001) Diagnostic performance of trauma US in identifying abdominal or pelvic free fluid and serious abdominal or pelvic injury. *Acad Radiol* 8:128-136.
7. Lee BC, Ormsby EL, McGahan JP, Melendres GM, Richards JR (2007) The utility of sonography for the triage of blunt abdominal trauma patients to exploratory laparotomy. *AJR Am J Roentgenol* 188:415-421.
8. Friese RS, Malekzadeh S, Shafi S, Gentilello LM, Starr A (2007) Abdominal ultrasound is an unreliable modality for the detection of hemoperitoneum in patients with pelvic fracture. *J Trauma* 63:97-102.
9. McGahan JP, Richards J, Fogata ML (2004) Emergency ultrasound in trauma patients. *Radiol Clin North Am* 42:417-425.
10. Abrams BJ, Sukumvanich P, Seibel R, Moscati R, Jehle D (1999) Ultrasound for the detection of intraperitoneal fluid: the role of Trendelenburg positioning. *Am J Emerg Med* 17:117-120.
11. Branney SW, Wolfe RE, Moore EE, Albert NP, Heinig M, Mestek M, Eule J (1995) Quantitative sensitivity of ultrasound in detecting free intraperitoneal fluid. *J Trauma* 39:375-380.
12. Shanmuganathan K, Mirvis SE, Sherbourne CD, Chiu WC, Rodriguez A (1999) Hemoperitoneum as the sole indicator of abdominal visceral injuries: a potential limitation of screening abdominal US for trauma. *Radiology* 212:423-430.